



A fNIRS investigation of event-related, Go-No-Go task in children

JongKwan Choi¹, Jae-Myoung Kim¹, Jaehyun Lim¹, Do-Joon Yi², Soyong Eom^{3*}

Optical Brain Electronics Laboratory, Seoul, South Korea¹;

Department of Psychology, Yonsei University, Seoul, Korea²; Epilepsy Research Institute, Yonsei University College of Medicine, Seoul, South Korea³



Objective

- Near-infrared spectroscopy (NIRS) provides a new direction for developmental cognitive neuroscience research.
- The application of fMRI to children is restricted to periods of tasks, however NIRS can be applied in the natural setting in children of any age, even whilst any task.
- The present study investigated functional brain activation patterns in children using functional Near Infrared Spectroscopy (fNIRS) during an event-related, Go-No-Go task and neuropsychological tests as a more cost-effective tool measuring cortical hemodynamics and a behavioral paradigm.

Neuropsychological Assessment	
Intelligence	Korean Wechsler Intelligence Scale -IV -Working Memory & Processing Speed
Executive function	Color Trails Test (CCTT)



Table 1. Neuropsychological assessment




Figure 2. Multi-distance based portable fNIRS device : NIRSIT

Methods

- This study enrolled 148 healthy normal participants (age range of 7.0–12.4 years old, 8.9 ± 1.5 years; 72 boys). Measurements were performed by NIRSIT-Lite (OBELAB Inc, Korea) which utilizes 5 LED sources and 7 photo detectors and it can measure 15 channels at prefrontal area.
- We adapted Go-No-Go task in an event-related version which might suggest executive processes such as interference resolution and response inhibition, and measured prefrontal activity during task performance.
- Relating neuropsychological tests of intelligence of Wechsler Intelligence Scale for Children –Fifth Ed. (WISC-V) including Verbal Comprehension, Visual Spatial, Fluid Reasoning, Working Memory and Processing Speed.
- The fNIRS data analysis was conducted on the concentration associated with the activation of each condition.
- Modified Beer-Lambert law (MBLL) is the main governing equation for NIRS, which extracts the concentrations of oxy- and deoxyhemoglobin by using light signals with two wavelengths.
- For representing individual distance matrix, density parameter is calculated after applying threshold value at the distance matrix.
- fNIRS density, GLM, and task parameters are calculated after applying threshold value at the distance matrix. fNIRS data and behavioral measure were analyzed with repeated measure ANOVA as a within-subjects and an inter-group factor. And correlation analysis was conducted among fNIRS, behavioral and neuropsychological data.




Figure 3. The Results of fNIRS density difference among neuropsychological tests on Go-No-Go task in children

Results

- In terms of development, no significant differences on fNIRS density, GLM, task parameters were shown, suggesting no prominent discrimination.
- However, the accuracy level of Go-No-Go test showed comparatively high correlation with density from fNIRS and showed high correlation with Processing Speed and Working Memory from WISC-V.

Conclusion

- Our results suggest that fNIRS parameters are associated with different functional activation patterns in the frontal subregions based on the characteristics of each task besides developmental aspects.
- fNIRS could be a potential measure as an effective tool to investigate the contributions of developmental prefrontal function of executive functioning in children.

Acknowledgment

The authors would like to thank the OBELAB for such generous support as providing us with NIRSIT device & technical consultation to the current study.

This research was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number:HI18C1166).

