

Rare events tend to be missed: can we predict behavioural errors using their neural signatures?

Hamid Karimi-Rouzbahani¹, Alexandra Woolgar^{*1,2}, Anina N. Rich^{*1}

¹ Perception in Action Research Centre and Department of Cognitive Science, Faculty of Medicine, Health, and Human Sciences, Macquarie University, Australia

² Medical Research Council Cognition and Brain Sciences Unit, University of Cambridge, UK

* Equally contributing authors

G100

Introduction

- When people monitor for rare targets, they are slower to respond and more likely to miss those targets especially in later stages of the task [1-3].

- There are many real-life situations where it can lead to tragic consequences such as in Ladbroke Grove Rail Crash, 1999.


- We asked, if we could detect what changes in the brain prior to a lapse in vigilance, could we prevent misses?

- To address this question, we built upon the extant literature of neuroimaging studies of vigilance [4] in three major ways:

* We designed a novel Multiple-Object Monitoring (MOM) paradigm presenting simultaneously moving objects simulating railway monitoring

* We used Magnetoencephalography (MEG), which provides high temporal resolution


* We used Multi-Variate Pattern Analyses (MVPA) and our recent "Error" data analyses [5], to predict behavioural errors



Methods

Multiple Object Monitoring (MOM) Paradigm

see video of an exemplar block here: https://osf.io/c6hy9/?view_only=ef87b9d629f843e99a96c63d2b942f52




MEG Setup




- Using MVPA, we extracted information about dots' distance from the deflection point (all possible pairs of 15 distances) and dots' direction of approach (right vs left).

Types of information decoded



Sample MEG data

- Band-passed filtered in 0.03-200 Hz



Bayes factor analysis was used for statistical tests


Statistical comparison	Null hypothesis	Alternative hypothesis
Decoding against chance	No difference	Difference exists
Decoding/RT/miss rate across conditions	No difference	Difference exists

Strong evidence for alternative: ● BF > 10
Moderate evidence for alternative: ○ 3 < BF < 10
Insufficient evidence for alternative: ○ 1 < BF < 3
Insufficient evidence for null: ○ 0.3 < BF < 1
Moderate evidence for null: ○ 0.1 < BF < 0.3
Strong evidence for null: ● BF < 0.1


Results

Behavioural and neural effects of vigilance decrement evoked by the MOM task


Deteriorating behavioural performance (i.e. increasing misses and reaction times) for the Monitoring vs Active condition shows that the MOM task evokes vigilance decrements successfully.



Decreased distance information in later blocks of the task (i.e. Time on task effect) and its interaction with Target frequency provides neural correlates for behavioural vigilance decrements: information decoding from brain activity decreases across time especially if relevant targets happen rarely.




Informational connectivity [6] decreased between Peri-occipital and Peri-frontal brain areas under Monitoring condition (vs Active) and when subjects missed (vs hit) the target. The weakened connectivity between sensory and cognitive areas may underlie vigilance decrements.



On miss (vs hit) trials:

Decoding of task-irrelevant information (i.e. direction of approach) is noisier, but not decreased. Decoding of task-relevant information (i.e. distance) is decreased.




Prediction of behavioural miss/correct trials using decoding of distance (task-relevant) information

The difference in information decoding rates between the correct vs miss trials allowed prediction of the outcome of the trial:

1. We trained 15 classifiers (per distance) using MEG data from correct trials and tested them using data from both correct and miss trials (A). The testing decoding rates were higher for correct vs miss trials (B).
2. In a second-level classification, we defined a threshold (B) that allowed the categorization of correct vs miss trials in the testing phase (C).
3. The threshold was calculated from the categorization rates of other subjects and was applied to the left-out subject.

- The results show accurate (~80%) prediction of miss/correct trials ~800 ms prior to the response time.



Summary

- Neural representation of task-relevant information decreases in brain activity with time on the task, especially when targets are infrequent.

- On miss trials, the decoding rates of task-relevant information decreases in the brain.

- We used these observations to predict behavioural outcome of individual trials with ~80% accuracy around 800 ms before the action was needed.

- These results provide new insights about how vigilance decrements are reflected in neural information decoding and offer an avenue for predicting behavioural errors using novel neuroimaging analyses.

References

- 1) Wolfe, J. M., Horowitz, T. S., & Kenner, N. M. (2005). Nature, 435(7041), 439-440.
- 2) Rich, A. N., Kunar, M. A., Van Wert, M. J., Hidalgo-Sotelo, B., Horowitz, T. S., & Wolfe, J. M. (2008). Journal of Vision, 8(15), 15-15.
- 3) Warm, J. S., Parasuraman, R., & Matthews, G. (2008). Human factors, 50(3), 433-441.
- 4) Langner, R., & Eickhoff, S. B. (2013). Psychological bulletin, 139(4), 870.
- 5) Woolgar, A., Dermody, N., Afshar, S., Williams, M. A., & Rich, A. N. (2019). bioRxiv, 673681.
- 6) Goddard, E., Carlson, T. A., Dermody, N., & Woolgar, A. (2016). Neuroimage, 128, 385-397.

Acknowledgement: This work was funded under the Australian Research Council (ARC)'s Discovery Projects funding scheme (DP170101780). A.W. was supported by an ARC Future Fellowship (FT170100105) and MRC intramural funding SUAG/052/G101400.