

Relationship of atrophy to task-related activity in the language network for different PPA clinical phenotypes

Rania J. Ezzo¹, Jeanne Gallée¹, Megan Quimby¹, Brad C. Dickerson¹, & Jessica Collins¹

¹Massachusetts General Hospital and Harvard Medical School, Boston, MA

MGH/HST Athinoula A. Martinos Center for Biomedical Imaging




Background

Primary Progressive Aphasia (PPA) is a neurodegenerative condition that is characterized by progressive language impairment. There are three subtypes, which generally have distinct clinical phenotypes. Semantic variant PPA (SvPPA) patients have most severe deficits in picture/object naming and single word comprehension. Nonfluent variant PPA (NfPPA) patients have dysfluent/effortful speech with errors in grammar comprehension. Logopenic variant PPA (LvPPA) patients are characterized by impairment in the auditory-verbal short-term memory known as the “phonological loop”¹²³.

Numerous studies have identified a network of brain regions that are selectively engaged during linguistic processing in the healthy brain. The subcomponents of this language network are differentially vulnerable to neurodegeneration in each of the distinct clinical phenotypes of PPA. The goal of this study was to investigate the effects of PPA on the functional integrity of the language network during linguistic processing.

Atrophy signatures

Cortical thickness of PPA compared to age-matched controls (n=48)



fMRI task design

Sentences Condition

THE BUTLER STARTED A RUMOR BUT THE COOK DID NOT BELIEVE HIM

Non-words Condition

GOR SCAVED CRE MISPRINE TICAITING UL IM LAS TROMER CRE CLUD CRE

Fedorenko Language Localizer⁴
Passive reading task
6 seconds per trial


Functional Analysis: methods & sample

- Single subject GLMs for each individual
- Functional ROIs defined for 6 language parcels per individual by selecting top 10% of voxels based on t-value
- fROIs used as a mask to determine mean Sentence > Nonwords contrast and mean cortical thickness z-score
- Interregional correlations computed by z-score of correlation matrix of spatially averaged BOLD signal per ROI


DIAGNOSIS	n	MEDIAN CDR (lang)	AGE RANGE	MEAN AGE	% MALE
SVPPA	8	1.0	63-70	66.7 yrs	38%
NfPPA	8	0.5	62-86	73.8 yrs	87%
LvPPA	11	0.75	60-79	69.5 yrs	64%
CN	8	n/a	32-61	48.7 yrs	13%

Defining functional regions-of-interest (fROIs)


- Angular Gyrus
- Posterior Temporal
- Anterior Temporal
- Middle Frontal Gyrus
- Inferior Frontal Gyrus
- Inferior Frontal Gyrus Pars orbitalis



Cortical thickness and functional activity in PPA




Cortical thickness GLM of whole PPA sample vs. controls



Whole brain GLM of sentences > non-words. The functional localizer activates the language network in the whole PPA sample. As expected, the activity is left lateralized and includes lateral temporal cortex, angular gyrus, inferior frontal gyrus, and caudal middle frontal gyrus.

Task-based functional connectivity



BOLD signal is significantly less correlated across regions in the LvPPA group compared to controls, with the language network fractionation being impacted most by the fractionation of the anterior temporal area.

Key Findings and Conclusions

Linguistic task-related activity is altered in PPA which is evident in the magnitude of functional activity compared to controls.

Our data demonstrates that the fedorenko localizer targets regions in the language network for the PPA sample, which warrants comparison to control group.

LvPPA patients, who have most prominent atrophy in the posterior temporal cortex and angular gyrus, have the most compromised functional connectivity across regions. This is consistent with the notion of this region serving as a hub for the language network.

References

- Gorno-Tempini, M. L., Hillis, A. E., Weintraub, S., Kertesz, A., Mendez, M., Cappa, S. F., ... & Manes, F. (2011). Classification of primary progressive aphasia and its variants. *Neurology*, 76(11), 1006-1014.
- Bonner, M. F., Ash, S., & Grossman, M. (2010). The new classification of primary progressive aphasia into semantic, logopenic, or nonfluent/agrammatic variants. *Current neurology and neuroscience reports*, 10(6), 484-490.
- Mesulam, M. M. (1987). Primary progressive aphasia—differentiation from Alzheimer's disease. *Annals of neurology*, 22(4), 533-534.
- Fedorenko, E., Hsieh, P. J., Nieto-Castañón, A., Whitfield-Gabrieli, S., & Kanwisher, N. (2010). New method for fMRI investigations of language: defining ROIs functionally in individual subjects. *Journal of neurophysiology*, 104(2), 1177-1194.