

Motivation

- **Hemispheric specialization (HS)** is the relationship between a cognitive, sensory, or motor function and a combination of brain structures of a given hemisphere.
- **Language lateralization and hand preference** are particular cases of HS [1, 2].
 - More than 90% of the overall population is right handed (RH) having their language function co-lateralized in the left hemisphere.
 - Around 20% of left handers (LH) present ambilateral or strongly-atypical language lateralization in the right hemisphere [3].
- Multiple studies of HS do not consider the linguistic competences of the participants, in particular, whether subjects speak one language (i.e. monolingual) or several languages (i.e. bilingual or multilingual) [4].
- Little is yet known about **how handedness affects brain lateralization in bilinguals**.

Experiment

- 111 early (age of acquisition < 6 years) highly-proficient Spanish-Basque bilinguals (mean age 26.19 ± 6.15 years; 59 Female)
- **MRI acquisition** in 3T Siemens Prisma^{Fit} MR scanner:
 - T1-w & T2-w anatomical images (1 mm³ isotropic)
 - 2 fMRI runs (GRE-EPI sequence, TR/TE=850/35 ms, multiband factor = 6, flip angle = 56°, voxel size = 2.4x2.4x2.4 mm³, 66 slices, matrix size = 88x88, 452 scans) while doing a **speech production task in Basque (BSQ) and Spanish (SPN)** [3].
- Subjects were presented during 1 sec white line drawing pictures on a black screen, which were either scrambled pieces randomly distributed or a cartoon depicting a scene. Afterwards, the subject had to covertly generate either the list of months of the year or a sentence describing the cartoon. More detailed information about the task in [3].




Figure 1. Illustration of the fMRI paradigm used for assessing language hemispheric lateralization.

Language lateralization in early bilinguals




Figure 2. Brain activation pattern of language production versus recitation. Brain activation pattern is shown for left (LH) and right handed (RH) bilinguals in both languages, Spanish (SPN) and Basque (BSQ). A paired t-test was performed to study the differences among languages (right part of the figure) and two sample t-test to study differences between LH and RH bilinguals (at the bottom).




Figure 4. Statistical analysis of LI scores. LEFT: Linear regression analysis of LI in Basque and Spanish for RH and LH and their corresponding correlation values. RIGHT: Box plots of the comparison between RH and LH for each language. Significant differences were found between RH vs LH bilinguals for each particular language, but not among languages.

- 1 Brain activation pattern of language production versus recitation is shown in **Figure 2**. Most active areas are part of the language processing network, such as inferior frontal gyrus ('Broca') and posterior temporal gyrus ('Wernicke'), precuneus, middle and superior temporal gyrus and frontal gyrus.
- 2 No significant differences were found after multiple comparison correction among languages, neither among left and right handed bilinguals when comparing language production versus recitation at the group level.
- 3 The distribution of LI scores shows more negative values in LH in comparison to RH, following previous findings [3]. Early bilinguals show more ambilateral activation in comparison to monolinguals (see **Figure 3**).




Figure 3. Distribution of LI scores in left and right handed bilinguals and for each language. Histograms with the distribution of the LI scores computed from the individual t-maps of sentence generation versus recitation contrast. A negative (positive) LI score means higher activation in the right (left) hemisphere. Values close to 0 means bilateral brain activation during the task.

- 4 Regression analysis show that some participants have different LI scores depending on the language, suggesting that we have to carefully consider the linguistic competences of each individual. A higher correlation value was found between the LI index of Spanish and Basque in LH in comparison to RH (see **Figure 4 - left**).
- 5 Significant differences were found in the LI scores of RH versus LH bilinguals in Spanish ($p=0.003$) and in Basque ($p=0.024$) (see **Figure 4 - right**).
- 6 No significant differences were found among languages in LH ($p=0.654$) and RH ($p=0.261$) (see **Figure 4 - right**).

Methods

- fMRI data was preprocessed in AFNI as follows: (i) slice time correction; (ii) head motion realignment; (iii) normalization to MNI template; and (iv) extraction of principal components from tissue-based WM and LV masks (based on Freesurfer) and edge-brain voxels [5, 6].
- The statistical parametric map of sentence generation versus recitation contrast was computed for each run in AFNI (3dREMLfit) including task-related and nuisance regressors.
- The hemispheric functional laterality index (LI) applied to the sentence generation versus recitation individual t-map was computed in SPM using the LI toolbox [7]:

$$LI = \frac{\sum \frac{activation_{left}}{mwf} - \sum activation_{right}}{\sum \frac{activation_{left}}{mwf} + \sum activation_{right}}$$

$mwf = \text{mask weighting factor}$

LI was computed with a bootstrap algorithm using the positive t-map, a lower bootstrap sample of 5 voxels and higher sample size of 1,000 voxels, and a resample ratio of $k=0.25$. Values ranging between -100 and +100, with -100 being a purely right and +100 a purely left activation.

- LI is computed for each participant and language (LI-SPN and LI-BSQ).
- Statistical inference was performed at (i) the group level to evaluate differences in the activation maps of the LH versus RH bilinguals (two sample t-test) and among languages (paired t-test); and (ii) the individual level to study the differences in LI scores of LH versus RH bilinguals.

Take-home message

- Early bilinguals show **more bilateral brain activation than monolinguals** in language-related areas during covert speech production.
- Our results call for the need to perform **analyses at the level of individual subjects** to investigate HS of language.
- Our study highlights the importance of **considering linguistic profiles** in determining HS in single subjects.

REFERENCES:

[1] Josse et al (2004), 'Hemispheric specialization for language', Brain Research Reviews, 44(1), 1-12.
 [2] Toga et al (2003), 'Mapping brain asymmetry', Nature Reviews Neuroscience, 4, 37-48.
 [3] Mazoyer et al (2014), 'Gaussian Mixture Modeling of Hemispheric Lateralization for Language in a Large Sample of Healthy Individuals Balanced for Handedness', PLOS ONE, 9(6), 1-14.
 [4] Hull et al (2007), 'Bilingual language lateralization: A meta-analytic tale of two hemispheres', Neuropsychologia, 45(9), 1987-2008.
 [5] Behzadi et al (2007), 'A component based correction method (CompCor) for BOLD and perfusion based fMRI', Neuroimage, 37(1), 90-101.
 [6] Patriat et al (2015), 'Using edge voxel information to improve motion regression for rs-fMRI connectivity studies', Brain connectivity, 5(9), 582-595.
 [7] Wilke et al (2007), 'LI tool: A new toolbox to assess lateralization in functional MR-data', Journal of Neuroscience Methods, 163(1), 128-136.

ACKNOWLEDGMENTS:

This research was supported by a fellowship from the Spanish Ministry of Science, Innovation and Universities (MICINN; FCI-2017-31814 to M. Termenon), La Caixa Foundation (ID 100010434, fellowship code LCF/BQ/IN17/11620063 to S. Moia), the Spanish Ministry of Economy and Competitiveness (Ramon y Cajal Fellowship, RYC-2017-21845 to C. Caballero-Gaudes), Beca Leonardo IN[18]_HMS_LIN_0058 (BBVA Foundation) to S. Mancini, the Spanish State Research Agency (BCBL "Severo Ochoa" excellence accreditation, SEV- 2015-490) and the Basque Government (BERC 2018-2021).