


INTRODUCTION

- Mindfulness-Based Interventions (MBIs) are a family of standardized cognitive and behavioral therapies that focus on cultivating mindfulness-related skills for improving maladaptive cognitive, emotional, and behavioral processes. MBIs have been developed for a wide range of clinical issues and populations in a variety of health settings.
- As the empirical evidence for the efficacy of these interventions continues to grow, the importance of investigating the mechanisms or processes by which they lead to beneficial outcomes is increasingly recognized
- The purpose of the present study was to investigate engagement between MBSR, perceived levels of stress and pain, and performance on an emotional variant of a Go-No-Go task


METHODS

- Pilot data (n = 9) was collected from a multi-site collaboration in individuals with moderate to high levels of stress
- Participants completed the Patient Reported Outcomes Measurement Information System (PROMIS-29) Pain Interference Scale and Perceived Stress Scale (PSS-10), followed by an emotional variant of a Go-No-Go task before and after MBSR
- Perceived levels of stress (PSS-10) and pain interference (PROMIS-29) were measured and analyzed
- Performance of the Go-No-Go task was measured by analyzing:
 - Errors of omission: incorrectly withholding a response to a Go trial
 - Errors of commission: incorrectly responding to a No-Go trial
 - A': A non-parametric index of sensitivity
 - Coefficient of Variation (CV): variability in reaction time




Example of Emotional Go-No-Go run


FIGURES




Error bars are SEM. * Significance $p = .041$




Error bars are SEM. * Significance $p = .037$




Error bars are SEM.




Error bars are SEM.




Error bars are SEM.




Error bars are SEM.




Error bars are SEM.




Error bars are SEM.




Error bars are SEM.




Error bars are SEM. * Significance $p = .020$




Error bars are SEM.



Error bars are SEM.



Error bars are SEM.



Error bars are SEM.

TABLES

Correlations Among Post-MBSR PROMIS Scores and Response Variability

	PROMIS	Negative CV	Positive CV	Neutral CV
PROMIS	1			
Negative CV	.791	1		
Positive CV	.927*	.959**	1	
Neutral CV	.913*	.933	.988**	1

**Correlation is significant at the .01 level (2-tailed)

*Correlation is significant at the .05 level (2-tailed)

RESULTS AND CONCLUSIONS

- Preliminary data showed a significant decrease in PSS and PROMIS scores
- Though not significant, there was an increase in errors of omission, and a decrease in errors of commission
- There was a slight increase in A' for negative and neutral words following MBSR, suggesting participants were better able to distinguish between Go trials and No-Go trials for negative and neutral words after completing MBSR
- There was a significant decrease in CV following MBSR, indicating participants were more consistent in reaction times after completing MBSR
- There was a significant correlation between post-MBSR PROMIS scores and post-MBSR response time variability, suggesting a relationship between perceived pain levels and engagement with the Go-No-Go task
- Taken together, these results suggest MBSR targets self-regulatory mechanisms, leading to changes in perceived stress and pain levels