

# Hebbian associative plasticity induced by a novel paired associative stimulation protocol shapes the properties of the Mirror Neuron System



Guidali Giacomo<sup>1,2</sup> & Bolognini Nadia<sup>1,3</sup>

<sup>1</sup> Department of Psychology, University of Milano-Bicocca & Milan Center for Neuroscience – NeuroMI, Milan, Italy.  
<sup>2</sup> PhD program in Neuroscience, School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy.  
<sup>3</sup> Laboratory of Neuropsychology, IRCCS Istituto Auxologico Italiano, Milan, Italy.



## 1 – BACKGROUND

The human brain is endowed with an action-observation network, the **Mirror Neuron System (MNS)** which implements a ‘mirror’ mechanism matching sensory and motor representations of actions. Even if the anatomo-functional properties of this network have been widely investigated, less is known about the plasticity mechanisms that rule mirror neurons and shape visuo-motor associations. One of the hypotheses put forward suggests that **mirror neurons develop their characteristics as a result of experience** and, in details, **Hebbian learning and Hebbian associative plasticity have been hypothesized as the neurophysiological substrate<sup>1</sup>**.

## 2 – AIM


Trying to deepen the nature of the plasticity mechanisms that rule the MNS, we test whether **atypical visuo-motor associations can be induced in the human MNS by directly targeting Hebbian associative plasticity using an ad-hoc developed non-invasive Paired Associative Stimulation (PAS) protocol<sup>2</sup>**, in turn re-shaping the resonance mechanisms of the human MNS.

## 3 – METHODS and MATERIALS

### 3.1 mirror PAS (m-PAS)


The m-PAS repeatedly paired (a) TMS pulses over the **right primary motor cortex** with (b) visual stimuli depicting a **right-hand index finger abduction movement**

**!! unilateral movements are not associated to motor resonance in the ipsilateral hemisphere<sup>3</sup>**




### 3.2 Action observation task

**Motor resonance** (i.e., effects of m-PAS) was assessed using a standard **action observation task<sup>4</sup>** divided in two blocks according to the side of the observed hand (left hand or right hand)



### 3.3 Experimental procedure

20 healthy participants tested in a two sessions within-subjects experiment




## 4 - RESULTS

\*  $p < 0.05$  \*\*  $p < 0.01$  \*\*\*  $p < 0.001$  (Bonferroni corrected)

### 4.1 Motor resonance before m-PAS


Preliminary, a rmANOVA was conducted to test whether **typical motor resonance phenomena is recorded before the administration of the m-PAS protocols**. A significant ‘viewed hand’ X ‘muscle’ interaction is found ( $F_{1,17} = 22.148$ ,  $p < .001$ ,  $\eta p^2 = .566$ )



MEPs facilitation in baseline only during the observation of left hands (ipsilateral to TMS) and only in the muscle involved in the observed movement (FDI)

### 4.2 Effects of m-PAS

m-PAS effects were assessed through a 2 “Muscle” X 3 “Condition” X 2 “viewed Hand” rm-ANOVA. A significant triple interaction was found ( $F_{2,34} = 4.31$ ,  $p = .021$ ,  $\eta p^2 = .202$ ) and it was further investigated in separate rmANOVAs for each muscle.



## 5 - CONCLUSIONS

The results of the present study show the efficacy of the m-PAS protocol, documenting that **it is possible to promote novel visuo-motor associations in the human MNS through the induction of plastic mechanisms that rely on Hebbian associative plasticity<sup>1</sup>**. Hebbian learning driven by the m-PAS is therefore a bottom-up, plastic process that starts with the **induction of associative plasticity only if we are exposed to visuo-motor association dealing with the time course of action execution (25 ms), rather than that of its visual input (250 ms)<sup>4</sup>**. Further studies have to be conducted to better explore, e.g., the role of other MNS cortical areas or the nature of the biological movements depicted (e.g., goal vs. non-goal movements; possible vs. impossible movements).

## 6 – REFERENCES

- 1) Keyser, C., & Gazzola, V. (2014). Hebbian learning and predictive mirror neurons for actions, sensations and emotions. *Philosophical Transactions of the Royal Society B*, 369, 20130175.
- 2) Suppa, A. et al. (2017). The associative brain at work: Evidence from paired associative stimulation studies in humans. *Clinical Neurophysiology*, 128(11), 2140–2164.
- 3) Aziz-Zadeh et al. (2002). Lateralization in motor facilitation during action observation: A TMS study. *Experimental Brain Research*, 144(1), 127–131
- 4) Naish, K. R. et al. (2014). Effects of action observation on corticospinal excitability: Muscle specificity, direction, and timing of the mirror response. *Neuropsychologia*, 64, 331–348.

CORRESPONDING AUTHOR:

[g.guidali@campus.unimib.it](mailto:g.guidali@campus.unimib.it)