Potential biomarker for ASD: Reduced pupil responses to repeated multisensory stimuli in young children with autism

¹ Autism and Neurodevelopmental Disorders Institute, ² Department of Psychological and Brain Sciences, ³ The George Washington University School of Nursing

Background

A failure to habituate to multi-sensory stimuli may underlie sensory hypersensitivities across disorders, e.g., Autism Spectrum Disorder (ASD); differences in sensory processing and habituation may be detectable early on and can serve as a biomarker for ASD¹.

Objectives

To investigate whether pupil responses can serve as a biomarker for multi-sensory processing and habituation, it is necessary to gain a mechanistic understanding of pupil responses and habituation thereof in neurotypical adults and children.


Sample & Design


Participants: 20 adults, 21 neurotypical children (M_{age} = 3.92yrs, SD = 1.2), 5 ASD children ($M_{age} = 4.17 yrs$, SD = 1.01)

Condition	Stimulus	Repetitions
Beeps	Optic flow field + beep/900ms	300 beeps (no ITI)
Bubblegum	5s animated gumball machine	50 trials (1-4s ITI)
Elf	10s jack-in-the-box movie clip	25 trials (2-8s ITI)

Data were collected via an Eyelink 1000 at 500Hz and preprocessed using the PsPM toolbox^{2,3.}

Pupil responses of neurotypical children and adults differ for higher complexity multisensory stimuli. Average Trial-Level Pupil Response





3 Kret, M. E and Sjak-Shie, E. E. (2018) Preprocessing pupil size data: Guidelines and code. *Behavior research methods, 51*(3).



Jonathan K Doyon^{1,2}, Ashley Darcey-Mahoney^{1,3}, Chynna Golding¹, Sarah Shomstein^{1,2}, & Gabriela Rosenblau^{1,2}

